






TABLE 1
THE SAMPLE OF OUTLYING CLUSTERS AND SURROUNDING FIELDS





Star Cluster	α_{1950} (h m s)	δ_{1950} ($^{\circ}$ ' ")	$l(^{\circ})$	$b(^{\circ})$	$r(^{\circ})$	$R(^{\circ})$
SL8,LW13	04 38 00	-69 07 37	280	-38.2	3.9	5.5
SL126,ESO85SC21	04 56 53	-62 36 37	272.5	-36.9	7.5	8.0
SL262,LW146	05 08 52	-62 26 27	272.0	-36.0	7.2	7.4
SL388,LW186	05 19 44	-63 31 44	273.1	-34.2	6.0	6.0
IC2134,SL437,LW198 ^a	05 24 47	-75 29 26	287.1	-31.7	6.0	6.0
SL451,LW206 ^a	05 25 57	-75 36 33	286	-31.6	6.1	6.1
SL509,LW221	05 29 29	-63 41 11	273.2	-33.1	5.9	6.0
SL817	06 01 09	-70 04 07	280	-29.7	3.4	4.8
ESO121SC03	06 01 27	-60 31 17	269.5	-29.4	10.3	11.4
SL842,LW399	06 07 53	-62 58 40	272.3	-28.8	8.4	9.9
SL862,LW431	06 14 04	-70 40 45	281.1	-28.6	4.5	6.3
OHSC33	06 16 33	-73 45 58	284	-28.5	5.7	6.9
OHSC37	07 08 01	-69 54 10	280	-24.0	9.2	13.0
An Inner Disk Cluster						
SL769 ^b	05 53 54	-70 04 44	281	-30.3	2.8	4.0





^aIC2134 and SL451 are located in the same frame.

^bSL769 is included for comparison purposes: its location corresponds to the inner disk, i.e. in our definition when the mean field turnoff becomes as bright as the clump. This occurs at $R \sim 4^{\circ}$ (see text).

















TABLE 3
AGES FOR CLUSTERS AND SURROUNDING FIELDS





Star Cluster	Age _{cluster} (Gyr)	Age _{field} (Gyr)
SL8,LW13	1.8:	2.0 ^c
SL126,ESO85SC21	2.2	>2.5
SL262,LW146	2.1	2.0
SL388,LW186	2.2	2.0 ^c
IC2134,SL437,LW198	1.0	2.0
SL451,LW206	2.2	2.0
SL509,LW221	1.2	1.5 ^c
SL817	1.5:	2.0 ^c
ESO121SC03	8.5	9.0
SL842,LW399	2.2:	2.5
SL862,LW431	1.8	2.0
OHSC33	1.4	2.0
OHSC37 ^a	2.1:	—
An Inner Disk Cluster		
SL769 ^b	1.8	—





^aThe field corresponds essentially to that of the Galaxy.

^bThe younger field turnoffs are younger than intermediate age (i.e. < 1Gyr) and outside of our age calibration.

^cA significant fraction of younger stars is present.














TABLE 4
 REDDENINGS AND METALLICITIES FOR CLUSTERS AND SURROUNDING FIELDS


Star Cluster	$E(B-V)_G$	$[Fe/H]_{cluster}$ this paper	$[Fe/H]_{cluster}$ Olszewski <i>et al.</i>	$[Fe/H]_{field}$
SL8,LW13	0.04	-0.55	—	-0.4
SL126,ESO85SC21	0.01	-0.5:	-1.18:	-0.5
SL262,LW146	0.00	-0.6	-0.34	-0.55:
SL388,LW186	0.03	-0.7	-0.76	-0.65
IC2134,SL437,LW198	0.10	—	—	-0.75
SL451,LW206	0.10	-0.75:	—	—
SL509,LW221	0.03	-0.9	—	-0.55
SL817	0.07	-0.55:	—	-0.75
ESO121SC03	0.03	-1.1	-0.93	—
SL842,LW399	0.03	-0.65	-0.36	-0.6:
SL862,LW431	0.09	-0.9	—	-0.65
OHSC33	0.09	-1.05:	-1.07:	-0.7
OHSC37	0.15	-0.7	-0.91	—
An Inner Disk Cluster				
SL769	0.08	-0.5	—	-0.55














Ages and Metallicities of Star Clusters and Surrounding Fields in the Outer Disk of the Large Magellanic Cloud

Eduardo Bica

Departamento de Astronomia, Instituto de Física, UFRGS, C.P. 15051, 91501-970

Porto Alegre RS, Brazil

Doug Geisler

Kitt Peak National Observatory, National Optical Astronomy Observatories,

P.O. Box 26732, Tucson, Arizona 85726

Horacio Dottori

Departamento de Astronomia, Instituto de Física, UFRGS, C.P. 15051, 91501-970

Porto Alegre RS, Brazil

Juan J. Clariá¹, Andrés E. Piatti¹

Observatorio Astronómico de Córdoba, Laprida 854, 5000,

Córdoba, Argentina

and

João F. C. Santos Jr.

Dep. de Física, ICEx, UFMG, C.P. 702, 30123-970 Belo Horizonte MG, Brazil

Received _____; accepted _____

Submitted to the Astronomical Journal

¹ Visiting Astronomer, Cerro Tololo Inter-American Observatory, which is operated by AURA, Inc., under cooperative agreement with the NSF.

ABSTRACT

We present Washington system C, T_1 color-magnitude diagrams of 13 star clusters and their surrounding fields which lie in the outer parts of the LMC disk ($r > 4^\circ$), as well as a comparison inner cluster. The total area covered is large ($2/3^\circ$), allowing us to study the clusters and their fields individually and in the context of the entire galaxy. Ages are determined by means of the magnitude difference δT_1 between the giant branch clump and the turnoff, while metallicities are derived from the location of the giant and subgiant branches as compared to fiducial star clusters. This yields a unique dataset in which ages and metallicities for both a significant sample of clusters and their fields are determined homogeneously. We find that in most cases the stellar population of each star cluster is quite similar to that of the field where it is embedded, thus sharing its mean age and metallicity. The old population ($t \geq 10$ Gyr) is detected in most fields as a small concentration of stars on the horizontal branch blueward and faintward of the prominent clump. Three particular fields present remarkable properties: (i) The so far unique cluster ESO121-SC03 at ≈ 9 Gyr has a surrounding field which shares the same properties, which in turn is also unique in the sense that such a dominant old field component is not present elsewhere, at least not significantly in the fields as yet studied. (ii) The field surrounding the far eastern intermediate age cluster OHSC 37 is noteworthy in the sense that we do not detect any evidence of LMC stars: it is essentially a Galactic foreground field. We can thus detect the LMC field out to $> 11^\circ$ (the deprojected distance of ESO121SC03), or ~ 11 kpc, but not to 13° (~ 13 kpc), despite the presence of clusters at this distance. (iii) In the northern part of the LMC disk the fields of SL388 and SL509 present color-magnitude diagrams with a secondary clump ≈ 0.45 mag fainter than the dominant intermediate age

clump, suggesting a stellar population component located behind the LMC disk at a distance comparable to that of the SMC. Possibly we are witnessing a depth effect in the LMC, and the size of the corresponding structure is comparable to the size of a dwarf galaxy. The unusual spatial location of the cluster OHSC37 and the anomalous properties of the SL 388 and SL 509 fields might be explained as debris from previous LMC interactions with the Galaxy and/or the SMC.

The mean metallicity derived for the intermediate age outer disk clusters is $\langle[\text{Fe}/\text{H}]\rangle = -0.7$ and for their surrounding fields $\langle[\text{Fe}/\text{H}]\rangle = -0.6$. These values are significantly lower than found by Olszewski *et al.* (1991, AJ, 101, 515) for a sample of clusters of similar age, but are in good agreement with several recent studies. A few clusters stand out in the age–metallicity relation in the sense that they are intermediate age clusters at relatively low metallicity ($[\text{Fe}/\text{H}] \approx -1$).

Subject headings: galaxies: Magellanic Clouds — galaxies: star clusters

1. INTRODUCTION

Unveiling the star formation history and chemical enrichment of galaxies is critical for understanding how they form and evolve. In this respect, Local Group galaxies play a fundamental rôle (Hodge 1989). The proximity of the Large Magellanic Cloud allows one to probe its stellar population properties with different techniques, namely photometry and spectroscopy of individual stars in clusters and the field, and integrated methods in the case of star clusters. Such studies for nearby galaxies are important to better understand very distant galaxies, whose stellar populations can only be probed by means of integrated properties.

Concerning field color-magnitude diagram (CMD) studies, ground-based observations have allowed the accurate study of the brighter evolutionary sequences, sampling relatively large fields throughout the LMC bar and disk (e.g. Butcher 1977, Hardy *et al.* 1984, Bertelli *et al.* 1992, Westerlund, Linde & Lyngå 1995, Vallenari *et al.* 1996). A prominent intermediate age (1-3 Gyr) stellar population is universally present in these fields, together with varying amounts of young blue main sequence (MS) stars. Hubble Space Telescope (HST) observations are limited to a small viewing area, but in turn allow deeper photometry, to well below the old MS turnoff. Several such fields have been studied with the V and I bands. One, at $\approx 4^\circ$ north of the bar (Gallagher *et al.* 1996) and another near the SE end of the bar (Elson, Gilmore & Santiago 1997) show evidence for the major star-forming event to have occurred ≈ 2 Gyr ago, in agreement with the ground-based studies. More recent HST studies (Holtzman *et al.* 1997, Geha *et al.* 1998) have investigated three LMC fields at 3° to 4° from the bar center, one located in the north-east and two in the north-west. Surprisingly, they are finding many more faint MS stars than expected and suggest that there has been more star formation in the past than previously believed. Their models, assuming a standard IMF slope, suggest that fully one half of the stars in these fields were

formed more than 4 Gyr ago. Although the latter studies refer to their fields as “outer”, we point out that in the present work we deal with genuine outer disk fields, well beyond any of these HST studies.

Integrated photometry of large star cluster samples of all ages have shown differences in the spatial distribution of age groups both in the bar region (Bica, Clariá & Dottori 1992) and the entire LMC (Bica *et al.* 1996). Differences in the spatial distribution among young groups have provided insight on the formation process and subsequent dynamical evolution of star cluster generations (Dottori *et al.* 1996). This integrated photometry cluster sample has been compared with integrated star cluster color models and has provided constraints on the cluster formation history (Girardi & Bica 1993, Girardi *et al.* 1995).

CMDs of LMC star clusters have also revealed a large intermediate age population (1-3 Gyr), which is separated by a pronounced age gap from the old stellar population as denoted by a few genuine globular clusters (see Da Costa 1991, Suntzeff *et al.* 1992, Olszewski, Suntzeff & Mateo 1996 for reviews). Recently, CMDs in the Washington system of a sample of candidate old clusters selected from the Bica *et al.* (1996) and Olszewski *et al.* (1991) studies revealed them to instead be of intermediate age (Geisler *et al.* 1997, hereafter Paper I). This study increased considerably the known sample of 1-3 Gyr old clusters with accurate age determinations, and reinforced the conclusion that a major formation epoch was preceded by a quiescent period of many Gyr, or that cluster dissipation has been more effective than generally believed (e.g. Olszewski 1993).

The objective of the present paper is to compare the properties of outer LMC clusters with those of their surrounding fields by using the same observational technique, and to infer the age-metallicity relation, and whether it depends on the spatial distribution throughout the LMC disk. In order to achieve this we employ Washington system C, T₁ bands and construct CMDs using the data of Paper I. Ages are inferred from the difference δT_1

between the giant branch clump and the turnoff (see also Paper I), and from the occurrence of particular stellar evolutionary sequences in the CMDs. The giant and subgiant branches allow one to derive metallicities using a technique analogous to that of Da Costa and Armandroff (1990) for VI photometry. However, our combination of Washington system filters is three times more metallicity sensitive than the VI system (Geisler & Sarajedini 1996, 1998), allowing us to obtain accurate metallicities for both the clusters and their fields. The cluster/field sample and the observations are described in Section 2. The cluster and field CMDs are described in Section 3. Ages and metallicities are derived in Section 4. In Section 5 we discuss the chemical enrichment of the outer disk, and in Section 6 the presence of dual clumps in two fields is noted and the possibility of a depth effect in this portion of the LMC disk is discussed. Finally, the conclusions of this work are given in Section 7.

2. The Sample and Observations

The outer LMC disk is tilted at $i \approx 45^\circ$ to the line of sight with the line of nodes at $\Theta \approx 7^\circ$, as indicated by the distribution of outlying star clusters (Lyngå & Westerlund 1963, see also Westerlund 1990 for a review). The total observed sample of clusters was first reported in Paper I. In the present paper we discuss in detail the 13 most distant outlying clusters from this sample and an additional inner cluster, SL769, included for comparison purposes, and their respective 13 surrounding fields (IC2134 and SL451 are located in the same frame). The star cluster designations in different catalogs and respective 1950 equatorial coordinates are listed in Table 1. Galactic coordinates are also given in the Table: since the distribution of the present clusters covers as much as 15° on the sky, variations of Galactic reddening are expected.

As a rule the clusters are centered in the frames, with some exceptions which were

shifted to avoid bright stars. In the case of IC2134 and SL451 the frame was centered approximately halfway between the two clusters, as illustrated in Figure 1, which also serves to illustrate typical clusters and field. The spatial distribution of the clusters is shown in Figure 2. With the exception of OHSC37 which is located far east away from the disk body, the remaining clusters are consistent with an inclined disk as described above. We also show in Table 1 the approximate projected angular distance from the bar center (taken as the position of the cluster NGC1928 $\alpha_{1950} = 5^h 21^m 19^s$, $\delta_{1950} = -69^\circ 31' 30''$), which in turn is $\approx 0.2^\circ$ south of the HI rotation curve center (see Westerlund 1990 for a review of centroids). Finally, the last column of Table 1 lists deprojected distances R assuming that all clusters are part of the inclined disk.

The observations were carried out with the CTIO 0.9m telescope in December, 1996 with the Tek2k #3 CCD, as described in Paper I. The scale on the chip is $0.40''$ per pixel, yielding an area $13.6' \times 13.6'$. SL769 was observed with the CTIO 4m in February, 1996 with the Tek2k #4 CCD, with similar pixel and areal coverage. The filters used for both runs were the Washington (Canterna 1976) C and Kron-Cousins R filters. The latter has significant throughput advantages over the standard Washington T_1 filter (Geisler 1996). In the present work, as in Paper I, we calibrate the observations in the C, T_1 system. In particular this filter combination allows us to derive accurate metallicities based on the standard giant branch technique outlined in Geisler & Sarajedini (1996, 1998). The data were reduced with the stand-alone version of DAOPHOT II (Stetson 1987) after trimming, bias subtraction and flat-fielding. More details on the observations, reductions and calibration procedures were given in Paper I.

After deriving the photometry for all detected objects in each filter, a cut was made on the basis of the parameters returned by DAOPHOT. Only objects with $\chi < 2$, photometric error $< 2\sigma$ more than the mean error at a given magnitude, and $|Sharp| < 0.5$ were kept in

each filter (typically discarding about 10% of the objects), and then the remaining objects in the C and T_1 lists were matched with a tolerance of 1 pixel, and raw photometry obtained. This raw photometry was then transformed to the standard Washington system as outlined in Paper I. Final calibrated photometry for each cluster (Table 2a-n) is available on the electronic AJ database.

We present in Figs. 3a to 3n the star cluster CMDs. The cluster radii were selected by eye judging the variation of the stellar density in the cluster surroundings, and ranged from 50-200 pixels, with a typical value of 75 pixels (30''). These radii are intended to optimize the cluster CMD sequences to avoid field contamination as much as possible. The radius in pixels for the extraction of each cluster is indicated in the corresponding CMD in Fig. 3.

We checked on the extent of field contamination of each cluster CMD by obtaining an equal-area field CMD composed of the addition of CMDs derived from 4 different fields, each of an area 1/4 that of the cluster and lying far away from the cluster. Note that such a comparison will overestimate field contamination since the photometric limit within the cluster will be brighter than in the field and more stars will be discarded from the cluster because of larger photometric errors, due to increased crowding. Nevertheless, in only 1 outlying cluster was the number of stars obtained in the equal-area field significantly more than 1/4 of the stars in the cluster CMD. The typical ratio was only $\sim 10\%$. Thus, we did not perform any statistical subtraction of the field contamination from the cluster CMDs.

We have also checked cluster extent on the basis of star counts and structural parameters and how it may be contributing to the surrounding fields in our frames. Three clusters in our sample have been studied by Kontizas, Hatzidimitriou & Kontizas (1987), namely SL 388 (LW 186), IC 2134 (LW 198) and SL 842 (LW 399) for which they could detect cluster stars as far as $r = 1.4'$, $r = 1.4'$ and $r = 1.7'$ respectively. Our selected radii for these clusters were $0.5'$, $0.5'$ and $0.4'$. Thus, our cluster radii are conservative and

should minimize field contamination.

We have also obtained CMDs for each field, excluding the pixels within a radius twice that used for the cluster, which minimizes contamination of the field by cluster stars. The tidal radii derived by Kontizas *et al.* for the three clusters ($r_t = 3'$, $r_t = 6.2'$ and $r_t = 3.1'$ respectively) are within the limits of the fields so defined. However, our criterion of field extraction is comparable to the detection limits of star counts, and the contribution of cluster stars to the field CMD, especially from clusters as relatively poorly populated as our sample, should not be significant. The tidal radius of ESO121-SC03 is $2.35'$ according to Mateo, Hodge & Schommer (1986). Notice that the cluster CMD extraction (Figure 3i) is for $r < 1.33'$, so that the field extraction corresponds to $r > 2.67'$, thus beyond the cluster tidal radius. The field CMDs are shown in Figs. 4a to 4m.

3. Description of CMDs

The cluster CMDs (Fig. 3) are typical of intermediate age clusters (IACs) with turnoffs ranging from magnitude levels slightly below the clump of He-burning stars to as much as 2 mag. below. The only exception is ESO121SC03 which is considerably older. See Paper I for a discussion of the clump to turnoff magnitude difference and its relation to age.

3.1. Fields

The uniform large size of the present frames is particularly suitable for the study of the brighter sequences of field stellar populations. The number of stars seen in the field CMDs (Fig. 4) is clearly correlated with the deprojected angular distances R from the bar center (Table 1). The radial dependance of several CMD features is also noteworthy. The field of OHSC37 at a distance $R=13^\circ$ has no evidence of a clump or horizontal branch or turnoff

of any kind, so that no LMC field stellar populations are detected in this frame. The wide distribution of stars in the diagram essentially consist of foreground Galactic stars. Note that OHSC37 is the most distant LMC cluster in the Olszewski *et al.* (1988) outer LMC cluster catalog. On the other hand, the field of ESO121SC03 presents clear clump and turnoff regions at the LMC distance. The $\delta T_1 \approx 2.9$ value is very similar to that of the cluster ESO121SC03 itself (Paper I), thus both cluster and LMC field have comparable ages (~ 9 Gyr) at this far north locus with $R=11.4^\circ$. No evidence exists for any intermediate age population in this field. One possibility is that the cluster and its surrounding field were a building block of the LMC, such as an accreted dwarf companion. Note that Galactic contamination is still important in this field.

Thus, we can detect the LMC field star population out to $> 11^\circ$ (the deprojected distance of ESO121SC03), or ~ 11 kpc, but not to 13° (~ 13 kpc), despite the presence of clusters at this distance. This is in good agreement with previous estimates of the extent of the LMC and its cluster system (e.g. de Vaucouleurs 1955, Lyngå and Westerlund 1963, Olszewski *et al.* 1988).

The LMC becomes prominent relative to the Galactic field at distances $10^\circ < R < 7^\circ$ (e.g. the SL126 and SL842 fields), and the principal turnoff is still well below the clump (≈ 2 mag). The oldest population is detectable by means of a concentration of stars near the instability strip locus faintward and blueward of the intermediate age clump.

For less distant fields the younger MS turnoff rapidly brightens, reaching the clump magnitude level for fields at $R \approx 5^\circ$ (e.g. that of SL8). The composite turnoff structure is obvious for such fields. Finally, the field of SL769 at $R=4^\circ$ definitely has an important component with a blue MS brighter than that of the clump. We adopt this criterion as a definition of the inner LMC disk. In this inner disk field, a minor old component, as demonstrated by the HB stars faintward and blueward of the clump, is still present.

The properties of our inner disk cluster sample and their fields will be the subject of a forthcoming paper.

4. Derivation of Ages and Metallicities

4.1. Ages

The utility of age determinations based on the magnitude difference between the clump/HB and the turnoff for IACs and old clusters is well known (see Phelps, Janes & Montgomery 1994 for Galactic open clusters). In Paper I we defined and calibrated such a method for δT_1 and applied it to our LMC cluster sample, including most of the present ones. In the present study we have rigorously eliminated photometric outliers. As a consequence, the CMDs were improved. We remeasured δT_1 values and in some cases small differences appeared with respect to Paper I values. Such differences were almost always within the errors, averaging only 0.15 Gyr. Age determinations by two independent investigators yielded ages within 0.3 Gyr in the mean, with a standard deviation of 0.3 Gyr. We also ranked the clusters according to turnoff morphology and reddening corrected magnitudes taking into account the foreground $E(B-V)_G$ values (Table 4). The derived ages are given in Table 3. These values should be preferred over those given in Paper I.

Since the fields are in most cases obviously composite in age, we measure the δT_1 value of the youngest well-populated turnoff and use the age calibration from Paper I to derive the representative age given in Table 3. Although there are some fields with small numbers of younger stars, most of the stars in these fields are as old as or older than the given age. We cannot place any useful constraints on the proportion of turnoff stars in each field older than ~ 4 Gyr as our photometry is not sufficient for the challenge. However, we do note that in the field with our deepest photometry (SL769, obtained with the 4m) one

can follow the subgiant branch down to a level which corresponds roughly with that shown by ESO121-SC03, i.e. the faintest subgiants in the SL769 field are consistent with an old (~ 9 Gyr) age. In some of the other fields, similarly faint subgiants can be seen.

Comparing the ages for the clusters with the surrounding fields as determined above we conclude that they are similar, which suggests that in general the cluster has the same origin as the surrounding field population.

4.2. Metallicities

Da Costa and Armandroff (1990) showed the utility of the $(V - I)$ color of the red giant branch for measuring metallicities in old stellar populations. This method now sees very wide use, and is the preferred technique, e.g., for HST WFPC2 observations of the stellar populations in distant Galactic globular clusters and nearby galaxies.

Geisler (1994) and Geisler and Sarajedini (1996) introduced a similar technique using the $(C - T_1)$ color of the Washington system and demonstrated that it had much potential for deriving metal abundances in distant objects, with a metallicity sensitivity greatly exceeding that of $(V - I)$. Geisler and Sarajedini (1998) have now derived the calibration of the standard giant branches in the Washington system technique. They have used the mean loci of giant and subgiant branches of Galactic globular and several old open clusters with known metallicities as fiducial clusters to derive the empirical relation between the $(C - T_1)$ color of the giant branch and metallicity, and show that this technique has three times the sensitivity to metallicity that the corresponding $(V - I)$ technique has. The ability to derive accurate metallicities for our program clusters and fields using this new technique was one of our primary motivations for using the Washington system in the current study.

However, the fiducial clusters used by Geisler and Sarajedini (1998) are all either typical Galactic globular clusters, with ages of > 10 Gyr, or among the oldest open clusters (M67, with an age of ~ 4 Gyr and NGC 6791, with an age of ~ 10 Gyr). In contrast, the only comparably aged cluster in the current sample is ESO121SC03 – all of the rest are IACs ranging from $\sim 1 - 2$ Gyr. Given the noticeable effect of such a large age difference on broadband colors, the Geisler and Sarajedini calibration is not directly applicable to our sample. Instead, what we have chosen to do is to use the Geisler and Sarajedini fiducial loci to derive metallicities for a sample of LMC and Galactic open clusters of intermediate age and well-determined metallicity and determine whether any offset is found between the known metallicity and the derived Washington value.

There are a total of five Galactic open clusters (Tombaugh 2, Melotte 71, NGC 2204, NGC 2506 and Melotte 66) and six LMC clusters (SL262, SL388, SL842, NGC 2213, OHSC33 and OHSC37) which have both good Washington photometry available for a number of stars along the upper giant branch (either this paper, previous publications or unpublished) and accurate metallicities (for the Galactic open clusters, these were taken from a variety of sources; for the LMC clusters, metallicities were taken from Olszewski *et al.* 1991). The ages of these clusters range from 1–4 Gyr.

In order to compare these clusters to the fiducial ones, a reddening and distance modulus were required to put the comparison clusters together with the standard ones in the $(M_{T_1}, (C - T_1)_0)$ plane. Again, for the Galactic open clusters we used the best current estimates available for these values based on an extensive literature search. For the LMC clusters, we assumed a true distance modulus $(m-M)_0=18.5$ taking into account results obtained from SN 1987A ($(m-M)_0=18.5$, Panagia *et al.* 1991) and the consequences of a recent revision of the Cepheid distance calibration on the LMC distance modulus ($(m-M)_0 = 18.50 \pm 0.15$, Madore & Freedman 1998). The extent of the LMC outer

disk is considerable, so we decided to use a foreground reddening $E(B-V)_G$ depending on the Galactic coordinates (Table 1) and the values from the maps by Burstein & Heiles (1982). The results are in Table 4. The reddening values for our entire LMC cluster sample vary from 0.00 to 0.10, except for OHSC37, which is located at a lower Galactic latitude and consequently has a higher reddening. Since we are dealing with the outer LMC, the disk internal reddening is expected to be negligible. We note that an increase of the assumed reddening by $E(B-V)=0.03$ decreases the derived metallicity (see below) by 0.12 dex.

We then derived metallicities for each of the 11 comparison clusters by interpolating by eye among the standard giant branches. Figure 5 shows the derived Washington metallicity vs. the standard value taken from the literature. A clear trend is found, indicating that the derived Washington metallicities for IACs using this technique require an approximately constant zero point correction. An unweighted mean yields a difference of 0.46 ± 0.21 dex. We then determined the metallicity of all of our program LMC clusters and fields in the same manner, and applied an offset of 0.45 dex to the derived value, i.e. we increased our estimate derived from a direct comparison with the standard clusters by 0.45 dex. Figure 6 shows a typical IAC and Figure 7 a typical IAC field. The final corrected values are given in Table 4. In a few cases, the metallicities were difficult or virtually impossible to determine because of the lack of bright giants – these cases are marked with colons or dashes in the Table and any derived metallicities are more uncertain. Note that the fields generally showed a significant range in metallicity, amounting to ~ 0.5 dex (although some of this scatter can be explained by Galactic field star contamination or LMC AGB stars), and that the values quoted are crude means. The offset is required for any intermediate age objects, which is the case for all but one of our LMC clusters and most of the field stars. However, note that our value for ESO121SC03, as derived from Figure 8, has not been corrected, since this is an old cluster. Our derived metallicity is only 0.17 dex lower than that of Olszewski *et al.* for this cluster, showing good agreement. In the

case of any field stars that are significantly older than ~ 4 Gyr, our corrected metallicity will represent an overestimate. However, such stars appear to be a very small minority in our fields.

In order to get an estimate of the errors involved in the metallicity derived from our interpolation procedure, two of us made independent measurements. We got the following differences: for metallicities of clusters and fields $\langle [\text{Fe}/\text{H}]_{EB} - [\text{Fe}/\text{H}]_{DG} \rangle = -0.05$ with $\sigma=0.07$; for the metallicity difference between cluster and respective field $\langle \Delta[\text{Fe}/\text{H}]_{EB} - \Delta[\text{Fe}/\text{H}]_{DG} \rangle = -0.07$ with $\sigma=0.13$ for 12 field/cluster CMDs. Therefore, we estimate that our internal metallicity errors are of the order of 0.1 dex and that the total metallicity uncertainties are ~ 0.2 dex.

Table 4 also lists the Olszewski *et al.* metallicity values for clusters in common. For these 7 clusters, we derive a mean difference of only 0.04 dex (our values are more metal-rich), with $\sigma = 0.34$ dex. Given that their errors are similar to ours, this is exactly the value expected if no other sources of error are present. Of course, we have used 5 of these clusters to derive the offset, but these represented $<1/2$ the total number of clusters used.

5. Chemical Enrichment

Olszewski *et al.* (1991) studied the chemical enrichment of the LMC based on ages from CMDs and metallicities derived from Ca II triplet spectroscopy of some individual giants. They obtained $\langle [\text{Fe}/\text{H}] \rangle = -0.42$ for 17 clusters with ages in the same range as for our sample (1–3 Gyr). In Paper I we increased the sample of well-studied IACs by combining ages from new CMDs with metallicities from Olszewski *et al.* (1991). We found evidence of a larger dispersion of metallicities at intermediate ages in the sense that some

outer clusters were more metal poor than the average. In the present study we revisit this issue for our outer disk sample and their fields, using the ages and metallicities derived here. Our method relies on a larger number of stars per cluster (Olszewski *et al.* generally observed only ~ 2 stars per cluster while we generally have 5 or more to determine the giant branch locus); the relative metallicity errors are similar.

The results are shown in Figure 9, where we plot clusters and fields, as well as some LMC globular clusters for comparison purposes (Paper I and references therein), and the mean curve for IACs derived by Olszewski *et al.* The resulting enrichment scenario we find for the outer disk is that of a pre-enriched gas with metallicity about one tenth solar (the metallicity level of ESO121SC03 in the quiescent epoch). Most parts of the outer disk were enriched by ≈ 0.3 dex before or during the burst which formed the IACs, but some regions apparently remained at the pre-enrichment level, forming some clusters with $[Fe/H] \approx -1.0$ even at later stages of the burst. Alternatively, such clusters might have been formed elsewhere and in their orbits are presently superimposed or embedded in slightly more metal rich outer disk fields (see discussions in Paper I). Indeed, Fig. 9 reveals a considerable range of metallicities for the IACs ($-0.5 > [Fe/H] > -1.05$), whereas the fields cover only 1/2 this range ($-0.4 > [Fe/H] > -0.75$). The clusters responsible for the metallicity dispersion are SL509, SL862 and OHSC33, which have ≈ 0.3 dex lower metallicities than the respective fields. Note that SL509 is not only metal-poor but also very young and that it exhibits the dual clump discussed in the next section. OHSC33 is also intriguing as another young, metal-poor cluster. The metallicity we derive for it is virtually identical to that found by Olszewski *et al.* However, for most of the sample, clusters and fields have comparable properties. The mean metallicity of the 11 IACs in the outer disk is $\langle [Fe/H] \rangle = -0.71 \pm 0.17$, while for the fields it is $\langle [Fe/H] \rangle = -0.61 \pm 0.11$. A direct comparison shows that the clusters are essentially all slightly more metal-poor than their respective fields, with a mean difference of 0.12 dex. Thus, the clusters and their

fields have very similar metallicities .

The IACs in the present sample are on the average 0.29 dex more metal poor than clusters of similar age in the Olszewski *et al.* (1991) sample. We have already shown that clusters in common yield similar metallicities. Probably our lower average metallicity is real and reflects a gradient between the outer and inner regions of the LMC (see also Paper I). Figure 10 plots metallicity vs. deprojected radius R for both the clusters and fields in our sample. No strong trend emerges but, as noted by Olszewski *et al.*, there is a tendency for the most distant clusters to be more metal-poor . Note that Olszewski *et al.* did uncover ~ 10 IACs with metallicities < -0.75 but their ages were mostly unavailable so they were not included in their age-metallicity relation.

Traditionally, the young ($< 1\text{Gyr}$) LMC population has been found to have a metallicity only slightly (~ 0.2 dex) less than solar (e.g. Harris 1983, Russell and Bessell 1989). IACs are generally regarded as having only slightly lower metallicities ($[\text{Fe}/\text{H}] \sim -0.4$), mostly based on the work of Olszewski *et al.* However, a number of studies have suggested lower metallicities for such objects might be more appropriate. In an early study including a large sample of IACs using narrow band integrated photometry, Bica, Dottori & Pastoriza (1986) derived metallicities similar to the present ones. Although ages for LMC IACs in that study were mostly based on LMC calibrators available at that time and require revision, the metallicities were derived from reference Galactic open and globular clusters with accurate abundances. Richtler *et al.* (1989) and Richtler (1993) derive values of -0.7 to -0.9 for several young LMC clusters using Strömgren photometry and high resolution spectroscopy. Vallenari *et al.* (1991) find a metallicity of -0.6 for the IAC NGC 2164. As for the LMC field, there are also several recent indications that we may need to revise metallicity estimates downwards. Geha *et al.* (1998) find very low metallicities ($\lesssim -1$) are required to fit the giant branch colors in their HST fields. And

Alves *et al.* (1998) analyze MACHO project photometry for $\sim 10^7$ stars in the bar and find a two-component model best fits the observations, with the intermediate age component having a metallicity of ~ -0.7 .

Thus, we conclude that intermediate age LMC clusters and fields, especially in the outer disk, may be more metal-poor than previously generally regarded. More spectroscopic and photometric studies with a large number of giants per cluster for a larger sample of IACs would help clarify this question.

6. Dual clumps

The fields of SL388 and SL509 (Fig.4), located together in the northern part of the LMC (Fig.2), present a unique feature: a populated secondary clump ≈ 0.45 mag fainter than the prominent main clump seen in all other fields. This fainter clump in the CMD is also slightly bluer, which suggests a lower metallicity and/or older age. The lower clump stars are found uniformly across both fields. This feature might be interpreted as a depth structure consisting of a layer of stars located approximately 10 Kpc behind the LMC, at a distance comparable to that of the SMC. The fact that the fields of SL509 and SL388 are separated by only $\approx 1^\circ$ on the sky sets a lower limit to the size of that possible structure of about 1 Kpc. An alternative explanation might be the presence of a dwarf companion to the LMC. In the SMC the occurrence of depth effects is well documented both in the field stellar component (Hatzidimitriou & Hawkins 1989) and in velocity space for the HI distribution (Mathewson & Ford 1984). A stellar bridge exists between the Clouds having arisen from their past interaction (Irwin, Demers & Kunkel 1990), and two concentrations in the bridge and one in the extreme of the SMC wing may evolve into dwarf galaxies (Bica & Schmitt 1995). All this evidence suggests that depth effects or dwarf galaxies might also occur elsewhere in the Magellanic System, as debris from interactions.

Arguments against this interpretation are that, if it is a superposition of two similar populations at different distances, one would expect at least greater scatter in the principal CMD sequences, yet this is not the case. More importantly, SL509 itself shows the same dual clump, with the fainter clump even more pronounced with respect to the brighter clump than seen in the field CMD, and an equal-area field reinforces this. Also, there is a hint of a dual clump in the very populous SL769 field, located many degrees away. More studies are needed to investigate this intriguing feature.

7. Conclusions

By using Washington system color-magnitude diagrams of a large sample of IACs and their surrounding fields in the LMC outer disk we were able to derive ages and metallicities. The cluster stellar population is a major component of the field where it is embedded, thus sharing its mean age and metallicity properties, except in three cases where the clusters are ≈ 0.3 dex more metal poor than the field, suggesting that the chemical enrichment was not globally homogeneous in the LMC. Our data is consistent with a scenario in which local star formation events generated both the clusters and a significant part of their surrounding stellar fields. Thus during the last 1-3 Gyr the dynamical evolution of the disk has not significantly taken them apart, which provides information on the diffusion time-scale and mixing of stellar generations in the disk. The old population ($t \geq 10$ Gyr) is detected in most fields as a small concentration of stars on the horizontal branch blueward and faintward of the prominent clump.

The unique cluster ESO121-SC03 at ≈ 9 Gyr has a surrounding field which shares the same stellar population. No other field so far studied is dominated by such an old population. One possibility would be that a field and cluster population coupling might last that long. Alternatively, we might be dealing with a building block recently accreted

by the LMC in the form of a dwarf galaxy.

One IAC cluster (OHSC37) is so far from the LMC body that no surrounding LMC field is detected. The present observations suggest that the LMC stellar disk extends out to between $\sim 11 - 13$ kpc in deprojected radius. In the northern part of the LMC outer disk the fields of SL388 and SL509 present evidence of a depth effect with a secondary component located behind the LMC disk at a distance comparable to that of the SMC. A background layer of stars in the LMC was possibly detected, and its size is at least ≈ 1 kpc, comparable to that of a dwarf galaxy. The peculiar location of the cluster OHSC37 and the depth effect in the SL388 and SL509 fields might be explained as debris from previous interactions of the LMC with the Galaxy and/or the SMC.

The average metallicity derived for the present outer disk IACs is $\langle [\text{Fe}/\text{H}] \rangle = -0.71 \pm 0.17$ and for their surrounding fields $\langle [\text{Fe}/\text{H}] \rangle = -0.61 \pm 0.11$. A few clusters stand out in the age–metallicity relation in the sense that they are intermediate age clusters at $[\text{Fe}/\text{H}] \approx -1$. The outer LMC disk (clusters and fields), at least of this age range (1-3 Gyrs) seem to be more metal-poor than previously generally regarded.

The authors would like to thank Cristina Torres for providing data critical to the metallicity calibration in advance of publication. E. Geisler, as always, was an inspiration. This research is supported in part by NASA through grant No. GO-06810.01-95A (to DG) from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. This work was partially supported by the Brazilian institutions CNPq and FINEP, the Argentine institutions CONICET and CONICOR, and the VITAE and Antorcha foundations.

REFERENCES

- Alves, D. *et al.* 1998, BAAS, in press
- Bertelli, G., Mateo, M., Chiosi, C. & Bressan, A. 1992, ApJ, 388, 400
- Bica, E., Clariá, J.J. & Dottori, H. 1992, AJ, 103, 1859
- Bica, E., Clariá, J.J., Dottori, H., Santos, J.F. Jr. & Piatti, A.E. 1996, ApJS, 102, 57
- Bica, E., Dottori, H. & Pastoriza, M. 1986 A&A, 156, 261
- Bica, E. & Schmitt, H.R. 1995, ApJS, 101, 41
- Burstein, D. & Heiles, C. 1982, AJ, 87, 1165
- Butcher, H. 1977, ApJ, 216, 327
- Canterna, R. 1976, AJ, 81, 228
- Da Costa, G.S. 1991, in The Magellanic Clouds, ed. R. Haynes & D. Milne (Dordrecht:Kluwer), 145
- Da Costa, G.S., & Armandroff, T.E. 1990, AJ, 100, 162
- de Vaucouleurs, G. 1955, ApJ, 60, 126
- Dottori, H., Bica, E., Clariá, J.J. & Puerari, I. 1996, ApJ, 742, 100
- Elson, R.A.W., Gilmore, G.F., Santiago, B.X. 1997, MNRAS, 289, 157
- Gallagher, J.S. *et al.* 1996, ApJ, 466, 732
- Geha, M.C. *et al.* 1998, preprint (astro-ph/9711144)
- Geisler, D. 1994, ESO Conf. Proc. No. 51, p.141
- Geisler, D. 1996, AJ, 111, 480
- Geisler, D. & Sarajedini, A. 1996, in Formation of the Galactic Halo...Inside and Out, ASP Conf. Ser., no.92, ed. H. Morrison & A. Sarajedini (San Francisco:ASP), 524

- Geisler, D. & Sarajedini, A. 1998, in preparation
- Geisler, D., Bica, E., Dottori, H., Clariá, J.J., Piatti, A.E. & Santos, J.F.C. Jr. 1997, AJ, 114, 1920
- Girardi, L. & Bica, E. 1993, A&A, 274, 279
- Girardi, L., Chiosi, C., Bertelli, G., & Bressan, A. 1995, A&A, 298, 87
- Hardy, E., Buonanno, R., Corsi, C.E., Janes, K.A. & Schommer, R.A. 1984, ApJ, 278, 592
- Harris, H.C. 1983, AJ, 88, 507
- Hatzidimitriou, D. & Hawkins, M.R.S. 1989, MNRAS, 241, 667
- Hodge, P.W., 1989, ARA&A, 27, 139
- Holtzman, J.A., *et al.* 1997, AJ, 113, 656
- Irwin, M.J., Demers, S. & Kunkel, W.E. 1990, AJ, 99, 191
- Kontizas, M., Hatzidimitriou, D., & Kontizas, E. 1987, A&AS 68, 493
- Lyngå G. & Westerlund, B.E. 1963, MNRAS, 127, 31
- Madore, B., & Freedman, W. 1998, ApJ 492, 110
- Mateo, M., Hodge, P. & Schommer, R.A. 1986, ApJ, 311, 113
- Mathewson, D.S. & Ford, V.L. 1984, IAU Symp. No. 108, 125
- Olszewski, E.W. 1993, in The Globular Cluster – Galaxy Connection, ASP Conf. Ser. No. 48, ed. G.H. Smith & J.P. Brodie (San Francisco:ASP), 351
- Olszewski, E.W., Harris, H.C., Schommer, R.A., & Canterna, R.W. 1988, AJ, 95, 84
- Olszewski, E.W., Schommer, R.A., Suntzeff, N.B., & Harris, H.C. 1991, AJ, 101, 515
- Olszewski, E.W., Suntzeff, N.B. & Mateo, M. 1996, ARA&A, 34, 511
- Panagia, N. *et al.* 1991, ApJ 380, L23

- Phelps, R. L., Janes, K.A. & Montgomery, K. A. 1994, AJ 107, 1079
- Richtler, T. 1993, in The Globular Cluster – Galaxy Connection, ASP Conf. Ser. No. 48, ed. G.H. Smith & J.P. Brodie (San Francisco:ASP), 375
- Richtler, T., Spite, M., & Spite, F. 1989, A&A, 225, 351
- Russell, & Bessell, M. 1989, ApJS, 70, 865
- Stetson, P.B. 1987, PASP, 99, 191
- Suntzeff, N.B., Schommer, R.A., Olszewski, E.W. & Walker, A.W. 1992, AJ, 104, 1743
- Vallenari, *et al.* 1991, A&AS, 87, 517
- Vallenari, A., Chiosi, C., Bertelli, G., Aparicio, A. & Ortolani, S. 1996, A&A, 309, 367
- Westerlund, B.E. 1990, A&A Rev., 2, 29
- Westerlund, B.E., Linde, P. & Lyngå G. 1995, A&A, 298, 39

Figure Captions

Fig. 1.— T_1 (R) frame of IC2134 (upper right cluster) and SL 451. The field is $13.6'$ on a side. North is up and east is to the left.

Fig. 2.— Spatial distribution of the sample. The LMC bar is schematically represented by a line for comparison purposes. Clusters in the outer disk are shown as triangles. The fields of SL388 and SL509 (filled triangles) present an extra clump in the CMD suggesting a depth structure (see Section 6). The field of SL769 (square) belongs to the inner disk.

Fig. 3.— Washington T_1 vs. $(C-T_1)$ CMDs of the star clusters. Extraction radius in pixels is given in each panel. a. SL8 b. SL126 c. SL262 d. SL388 e. IC2134 f. SL451 g. SL509 h. SL817 i. ESO121SC03 j. SL842 k. SL862 l. OHSC33 m. OHSC37 n. SL769

Fig. 4.— Washington T_1 vs. $(C-T_1)$ CMDs of the surrounding fields, excluding areas of radius two times that of the cluster. a. SL8 b. SL126 c. SL262 d. SL388 e. IC2134 f. SL509 g. SL817 h. ESO121SC03 i. SL842 j. SL862 k. OHSC33 l. OHSC37 m. SL769

Fig. 5.— Metallicity derived from the color of the giant branch in the Washington system compared to standard giant branches vs. standard metallicity. Plus signs are LMC clusters, squares are Galactic open clusters. The solid line shows perfect correlation; the dashed line indicates the mean relation we find: $[Fe/H]_{STD} = [Fe/H]_{SGB} + 0.45$.

Fig. 6.— Metallicity derivation for the IAC SL862. The cluster has been placed in the absolute T_1 magnitude – dereddened $(C - T_1)$ color plane assuming a true distance modulus of 18.5 and a reddening of $E(B-V)=0.09$. The standard giant branches are those of NGC 6752 (offset $[Fe/H]=-1.1$) and NGC 1851 (offset $[Fe/H]=-0.75$).

Fig. 7.— Metallicity derivation for the field of SL862. Caption as for Figure 6, with the addition of the standard giant branch for 47 Tuc (offset $[Fe/H]=-0.3$).

Fig. 8.— Metallicity derivation for the old cluster ESO121-SC03. Caption as for Figure 7, except that the standard, not offset, metallicities are shown.

Fig. 9.— Chemical evolution of the LMC outer disk: open triangles refer to fields, filled triangles to clusters. Lines link cluster and the respective surrounding field. For comparison purposes we show some LMC globular clusters (filled circles) and an LMC inner disk cluster (open square). The age for the field corresponding to SL126 located at (9.4, -0.5) is a lower limit (Table 3). The curve shows the mean age-metallicity relation derived by Olszewski *et al.* (1991) for IACs.

Fig. 10.— Metallicity vs. deprojected radius for LMC outer clusters (squares) and fields (plus signs). The point at the bottom right is ESO121-SC03.